

Deceleration and Trapping of Cold Free Radicals by Pulsed Magnetic Fields

Towards Ultra-high Resolution Spectroscopy and Cold Chemistry



Canadian Centre
for Research on
Ultra-Cold Systems



Canada Foundation
for Innovation

Fondation canadienne
pour l'innovation

Takamasa Momose
The University of British Columbia
Centre for Research on Ultra-Cold Systems (CRUCS)
Chemistry / Physics and Astronomy
Vancouver, CANADA

Cold Molecules

Molecules with a low translational energy

Cold Molecules

Molecules with a low translational energy

1. Ultra-high Resolution/Precision Spectroscopy

Molecules with a low translational energy

1. Ultra-high Resolution/Precision Spectroscopy

The Origin of spectral linewidth

Natural $\frac{1}{2\pi\tau_{sp}}$

Doppler $\sqrt{\pi \ln 2} \tilde{\nu} \langle v \rangle$

Transit time $\frac{1}{2\sqrt{2 \ln 2}} \frac{\langle v \rangle}{d}$

Cold Molecules

Molecules with a low translational energy

1. Ultra-high Resolution/Precision Spectroscopy

The Origin of spectral linewidth

$$\begin{aligned}
 \text{Natural} & \quad \frac{1}{2\pi\tau_{sp}} \\
 \text{Doppler} & \quad \sqrt{\pi \ln 2} \, \tilde{\nu} \langle v \rangle \\
 \text{Transit time} & \quad \frac{1}{2\sqrt{2 \ln 2}} \frac{\langle v \rangle}{d}
 \end{aligned}$$


CH₃ (*M* = 15) *v*₃ (3,160 cm⁻¹) *μ*₁₀=0.1 D

<i>T</i>	Mean velocity < <i>v</i> > [m/s]	Doppler [MHz]	Transit time* [kHz]	Natural [Hz]
300 K	650	303	276	16
1 K	38	17.5	16	16
1 mK	1.2	0.55	0.5	16

***1 mm diameter laser radiation**


High-Resolution Spectroscopy

ν_3 Absorption of CH_3 (slit jet)



Davis et al. J. Chem. Phys. 107, 5661 (1997)

solid line. The least-squares fits indicate an OH Doppler width of $\Delta\nu = 105(2)$ MHz, which is in reasonable agreement with $\Delta\nu = 98(3)$ MHz predictions from scaling the experimentally observed HF dimer Doppler widths by the ratio of absorption frequencies and the square root of the mass. This would be consistent with a slit translational temperature of ≈ 30 K and \approx fourfold enhancement in spectral resolution over Doppler limited linewidths under typical discharge cell conditions.



30 K ~ 200 m s⁻¹

Cold Molecules

Molecules with a low translational energy

1. Ultra-high Resolution/Precision Spectroscopy

The Origin of spectral linewidth

Natural $\frac{1}{2\pi\tau_{sp}}$


Doppler $\sqrt{\pi \ln 2} \tilde{\nu} \langle v \rangle$

Transit time $\frac{1}{2\sqrt{2 \ln 2}} \frac{\langle v \rangle}{d}$

CH₃ (*M* = 15) *v*₃ (3,160 cm⁻¹) *μ*₁₀=0.1 D

<i>T</i>	Mean velocity <i><v></i> [m/s]	Doppler [MHz]	Transit time* [kHz]	Natural [Hz]
300 K	650	303	276	16
1 K	38	17.5	16	16
1 mK	1.2	0.55	0.5	16


***1 mm diameter laser radiation**



Ultrahigh-resolution spectroscopy
Test of fundamental symmetries
Constancy of fundamental constants
eEDM measurement


2. Collision Dynamics / Coherent Control

Relevant length: Thermal de Broglie length




2. Collision Dynamics / Coherent Control


Relevant length: Thermal de Broglie length




matter wave



thermal fluctuation of momentum




superposition of matter wave




2. Collision Dynamics / Coherent Control


Relevant length: Thermal de Broglie length




matter wave



thermal fluctuation of momentum



superposition of matter wave



thermal de Broglie length

	H ₂	CH ₄	C ₆ H ₆
M(amu)	2	16	78
300K	70 pm	25 pm	11 pm
1 K	1.3 nm	440 pm	200 pm
0.1 mK	130 nm	44 nm	20 nm
10 nK	13 μm	4.4 μm	2 μm
"classical" size	1 Å 100 pm	3 Å 300 pm	5 Å 500 pm


large matter wave : below 1 K

Cold Ion-Molecule Reactions


Tim Softley's Group

<http://physchem.ox.ac.uk/~tps/iquad.htm>

Cold ion-molecule reactions



Negative temperature effect
in ion-neutral reaction




Cold Ion-Molecule Reactions

Tim Softley's Group

<http://physchem.ox.ac.uk/~tps/iquad.htm>

Cold ion-molecule reactions




Negative temperature effect in ion-neutral reaction

Interstellar Chemistry


T: 5K - 100 K

neutral-neutral reactions?




2. Collision Dynamics / Coherent Control


Relevant length: Thermal de Broglie length




matter wave



thermal fluctuation of momentum



superposition of matter wave



thermal de Broglie length

	H ₂	CH ₄	C ₆ H ₆
M(amu)	2	16	78
300K	70 pm	25 pm	11 pm
1 K	1.3 nm	440 pm	200 pm
0.1 mK	130 nm	44 nm	20 nm
10 nK	13 μm	4.4 μm	2 μm
"classical" size	1 Å 100 pm	3 Å 300 pm	5 Å 500 pm

large matter wave : below 1 K

Collision and reaction dynamics
Interference of molecular dB waves
Coherent control of reactions
Separation of chiral molecules
Quantum information processes


Motivation

Spatially confine (chemically unstable) molecules

Ultra-high-resolution spectroscopy
Cold Chemistry



DC static fields
Paul trap
Anti-Helmholtz trap



AC fields
Optical lattice
MW trap


translational velocity : $v < 10 \text{ m s}^{-1}$

Generation of Translationally Cold Molecules


Buffer gas cooling

Doyle (Harvard)

He ~ 1 K
Target molecules



thermalization




Counter rotating source

- Mechanical deceleration

Herschbach (Harvard)

Stienkemeier, Mudrich (Freiburg)


Momose (UBC)



Stark effect on polar molecules

- Stark deceleration


Meijer (Max-Planck Institute) etc



Generation of Translationally Cold Molecules

Buffer gas cooling

Doyle (Harvard)





Counter rotating source

- Mechanical deceleration

Herschbach (Harvard)

Stienkemeier, Mudrich (Freiburg)


Momose (UBC)




Stark effect on polar molecules

- Stark deceleration

Meijer (Max-Planck Institute) etc





Zeeman Decelerator



H atoms F. Merkt et al. PRA **76**, 023412 (2007).
 Ne atoms MG Raizen, PRL **100**, 090303 (2008).


Zeeman Decelerator




H atoms F. Merkt et al. PRA **76**, 023412 (2007).
 Ne atoms MG Raizen, PRL **100**, 090303 (2008).

Zeeman Decelerator


A series of Solenoid coils



Solenoid coil




Zeeman splitting of CH_3




H atoms F. Merkt et al. PRA **76**, 023412 (2007).
 Ne atoms MG Raizen, PRL **100**, 090303 (2008).

Zeeman Decelerator

A series of Solenoid coils




Solenoid coil



Zeeman splitting of CH_3


$|N, |K|\rangle$




H atoms F. Merkt et al. PRA **76**, 023412 (2007).
 Ne atoms MG Raizen, PRL **100**, 090303 (2008).

Zeeman Decelerator


A series of Solenoid coils



Solenoid coil





Zeeman splitting of CH_3




H atoms F. Merkt et al. PRA **76**, 023412 (2007).
 Ne atoms MG Raizen, PRL **100**, 090303 (2008).

Stark Decelerator





Meijer et al,
Phys. Rev. A **65**, 53416 (2002)


Stark splitting of CH_3I



Stark Decelerator





Stark splitting of CH_3I




Meijer et al,
Phys. Rev. A **65**, 53416 (2002)

Stark Decelerator



Meijer et al,
Phys. Rev. A **65**, 53416 (2002)

Stark splitting of CH_3I



Zeeman vs Stark Effects

	Stark	Zeeman
Hamiltonian	$\hat{H}_S = -\mu \cdot \mathbf{E}$	$\hat{H}_Z = -\mu \cdot \mathbf{H}$
dipole moment	electric dipole 1 D = 3.3564×10^{-30} Cm	magnetic dipole 1 $\mu_B = 9.274 \times 10^{-24}$ JT ⁻¹
field	electric 20 kV / 5 mm = 4 MV m ⁻¹	magnetic 10 T
shift	$\mu E = 9.72 \times 10^{-5}$ aJ = 4.9 cm ⁻¹ = 7.0 K	$\mu_B H = 9.28 \times 10^{-5}$ aJ = 4.6 cm ⁻¹ = 6.8 K

Magnetic Field

Free radicals have a large permanent magnetic moment due to an unpaired electron.

Hamiltonian

$$\hat{H} = \hat{H}_R + \hat{H}_{SR} + \hat{H}_{hf} + \hat{H}_Z$$

Zeeman term


$$\hat{H}_Z = \mu (g_\ell \mathbf{L} + g_S \mathbf{S} + g_I \mathbf{I} + g_N \mathbf{N}) \cdot \mathbf{H}$$

At high field, spin is well decoupled

$$|NM_NK\rangle |SM_S\rangle$$

Boland et al.
Proc. R. Soc. Lond. A 360, 507 (1978).

HCO Zeeman splitting



Zeeman Decelerator vs Stark Decelerator


$$|NM_N K\rangle |SM_S\rangle$$

Only LFS are decelerated


Zeeman splitting of CH_3

$|N, |K|\rangle$

M_S



Stark shift/splitting of CH_3I



Zeeman Decelerator vs Stark Decelerator


$$|NM_N K\rangle |SM_S\rangle$$

Only LFS are decelerated


Zeeman splitting of CH₃

$|N, |K|\rangle$

M_S



Stark shift/splitting of CH₃I



Lower rotational states become HFS at high fields

The rotational ground state cannot be decelerated.

Only small molecules

Zeeman Decelerator vs Stark Decelerator


$$|NM_N K\rangle |SM_S\rangle$$

Only LFS are decelerated


Zeeman splitting of CH₃

$|N, |K|\rangle$

M_S



Stark shift/splitting of CH₃I



Any ground rotational state has a LFS component.

The ground rotational state can be decelerated.

More universal

Lower rotational states become HFS at high fields






The rotational ground state cannot be decelerated.

Only small molecules

High Field Solenoid Coil @ CRUCS

30 turns Solenoid coil



M. Raizen (Texas, Austin)



High Field Solenoid Coil @ CRUCS

30 turns Solenoid coil

M. Raizen (Texas, Austin)





$$v=400 \text{ m/s} \rightarrow 4\mu\text{s}/1.5 \text{ mm}$$


High Field Solenoid Coil @ CRUCS

30 turns Solenoid coil


M. Raizen (Texas, Austin)



$$v=400 \text{ m/s} \rightarrow 4\mu\text{s}/1.5 \text{ mm}$$





1st Generation Zeeman Decelerator



Phys. Chem. Chem. Phys **15**, 1772 (2013)

1st Generation Zeeman Decelerator



Phys. Chem. Chem. Phys **15**, 1772 (2013)



1st Generation Zeeman Decelerator

Liq. N₂ cooling


total resistance

0.42 Ω (RT)

0.36 Ω (liq N₂)




SPECIFIED




700A : 4.9 T

Test Sample : O₂



$$\hat{H} = \hat{H}_R + \hat{H}_{SR} + \hat{H}_{SS} + \hat{H}_Z$$



Simulation

$$^{16}\text{O}_2 \quad |N, J, M_J\rangle = |1, 2, 1\rangle$$

initial velocity : 250 m/s
400 A switching

transverse distribution

molecular
distribution

velocity (m/s)

340
320
300
280
260
240
220
200
180
160

longitudinal
distribution

coil switching

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15

15 coils

Simulation


$$^{16}\text{O}_2 \quad |N, J, M_J\rangle = |1, 2, 1\rangle$$

initial velocity : 250 m/s
400 A switching


transverse distribution

molecular
distribution


velocity (m/s)



longitudinal
distribution



coil switching



1 2 3 4 5 6 7 8 9 10 11 12 13 14 15

15 coils

Simulation

$$^{16}\text{O}_2 \quad |N, J, M_J\rangle = |1, 2, 1\rangle$$

initial velocity : 250 m/s
400 A switching

transverse distribution

molecular
distribution

velocity (m/s)

340
320
300
280
260
240
220
200
180
160




longitudinal
distribution

coil switching

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15


15 coils

O₂ Detection : 2+1 REMPI



McCann, Chen, Datskou, Evans, Chem Phys. **174**, 417 (1993)

O₂ REMPI Detection





assignments

Katsumata et al.



J. Elec. Spec. Rel. Phenom. **41**, 325 (1986).

Zeeman Deceleration of O₂




Zeeman Deceleration of O₂

Simulation




Zeeman shift





Zeeman Deceleration of O₂


Experiment




Simulation




Zeeman shift




2nd Generation : 80 Coils



2nd Generation : 80 Coils




“CRUCS” Pulsed Valve



Opening time $\sim 25 \mu\text{s}$



Low & High stagnation pressure (1 - 10 bar)
able to operate at liq. He temp



Deceleration of O₂

80-coil system



O₂ seeded in Kr @ 135 K (65 us open)



Deceleration of O₂

80-coil system



O₂ seeded in Kr @ 135 K (65 us open)



Deceleration of O₂




80-coil system

O₂ seeded in Kr @ 135 K (65 us open)




REMPI Spectra of Decelerated O₂

$|J=1, M_J=0,1\rangle$ (and $|J=2, M_J=2\rangle$) are decelerated.




Trapping of Decelerated O_2

Anti-Helmholtz Coil





Trapping of Decelerated O_2

Anti-Helmholtz Coil



Trapping of O₂ in a Static Magnetic Field

anti-Helmholtz coil trap







Trapping of O₂ in a Static Magnetic Field


anti-Helmholtz coil trap

600 A


$E_{\perp} \sim 1\text{T} \sim 1.1\text{cm}^{-1}$
 $\sim 28\text{m/s (O}_2\text{)}$





$E_{\parallel} \sim 4\text{T} \sim 3.6\text{cm}^{-1}$
 $\sim 52\text{m/s (O}_2\text{)}$






Trapping of O₂ in a Static Magnetic Field





Trapping of O₂ in a Static Magnetic Field



Trapping of O₂ in a Static Magnetic Field





Trapping of O₂ in a Static Magnetic Field





simulation

Trap ON

Trap OFF



Trapping of O₂ in a Static Magnetic Field





simulation

Trap ON




Trap OFF

**The first trapping signal
of Zeeman decelerated molecules**



REMPI Spectra of Decelerated O₂


O₂ in the $|J=1, M_J=0,1\rangle$ state are trapped



Deceleration of Free radicals

Deceleration of Free radicals

Methyl Radical: CH₃



High-resolution Spectroscopy

A planer oblate top molecule

First Identification:

G. Herzberg and J. Shoosmith, Can. J. Phys. **34**, 523 (1956)

Matrix isolation spectroscopy:

D. E. Milligan and M. E. Jacox, J. Chem. Phys. **47**, 5146 (1967)

Flash Photolysis IR spectroscopy:

L. Y. Tan, A. M. Winer, and G. C. Pimentel, J. Chem. Phys. **57**, 4028 (1972).

v₂ band high-resolution IR (606.45 cm⁻¹):

C. Yamada, E. Hirota, and K. Kawaguchi, J. Chem. Phys. **75**, 5256 (1981)

C. Yamada, and E. Hirota, J. Chem. Phys. **78**, 669 (1983)

$$\mu_{10}=0.28 \text{ D}$$

v₃ band high-resolution IR (3160.82 cm⁻¹):

T. Amano, P. Bernath, C. Yamada, Y. Endo, and E. Hirota,
J. Chem. Phys. **77**, 5284 (1982)


$$\mu_{10}=0.1 \text{ D}$$

(slit jet) S. Davis, D.T. Anderson, G Duxbury, and D. J. Nesbitt,
J. Chem. Phys. **107**, 5661 (1997)

Detected in Space


H. Feuchtgruber, F.P. Helmich, E.F. van Dishoeck and C.M. Wright,
Astro. J. **535**, L111 (2000).

Reactive Species




Methane oxidation mechanism

Reactive Species




Methane oxidation mechanism

CH₃ + OH Barrierless reaction





Ahren W. Jasper et al,
J.Phys.Chem. A **111**, 3932(2007)

Reactive Species




Methane oxidation mechanism




A. M. Morrison et al,
J.Phys.Chem.A **116**, 5299(2012)

CH₃ + OH Barrierless reaction



Ahren W. Jasper et al,
J.Phys.Chem. A **111**, 3932(2007)

CH₃ + OH Barrierless reaction




Ahren W. Jasper et al,
J.Phys.Chem. A **111**, 3932(2007)



A. M. Morrison et al,
J.Phys.Chem.A **116**, 5299(2012)

Cold chemistry !

ortho-para modification



Davis et al. J. Chem. Phys. 107, 5661 (1997)

study of ortho-para conversion

Closed cycle IR transitions

IR laser cooling

Sympathetic cooling with Li/Rb

T.V. Tscherbul et al.
Phys. Rev. Lett. **106**, 073201 (2011).

Various possibilities to cool
below 1 mK



Production of CH_3

CRUCS valve

Liq. N_2 cooling

Opening time : 30~60 μs

4- 6 bar pressure





Discharge period: 150 μs
FWHM beam width 50 μs

A compact radical beam

CH₃ Detection : 2+1 REMPI

B. Martínez-Haya, et al. Chem. Phys. Lett. 311, 159 (1999)





Rydberg transition

resonant at $4 \times 17,470 \text{ cm}^{-1}$

O₂ at $4 \times 17,400 \text{ cm}^{-1}$

CH₃ Detection : 2+1 REMPI

B. Martínez-Haya, et al. Chem. Phys. Lett. 311, 159 (1999)




Rydberg transition

resonant at $4 \times 17,470 \text{ cm}^{-1}$

O₂ at $4 \times 17,400 \text{ cm}^{-1}$


Black, Powis J. Chem. Phys. 89, 3986 (1988)




rotational assignments

CH₃ Radical Beam

Black, Powis J. Chem. Phys. 89, 3986 (1988)



**Discharge of CH₄/Ar
with a cold nozzle (150K)**



rotational temp <5 K

N''=0 : N''=1 2:1

CH₃ Radical Beam

Detection is 1 m downstream of a nozzle.

Requirements for the Zeeman deceleration:

High-density, low rotational temperature, well collimated, slow,
and background free radical beam

CH₃ Radical Beam

Detection is 1 m downstream of a nozzle.

Requirements for the Zeeman deceleration:

High-density, low rotational temperature, well collimated, slow,
and background free radical beam

UV dissociation of CH₃I

CH₃ Radical Beam

Detection is 1 m downstream of a nozzle.

Requirements for the Zeeman deceleration:

High-density, low rotational temperature, well collimated, slow,
and background free radical beam

UV dissociation of CH₃I

UV dissociation or discharge of Di-tert-butyl peroxide

CH₃ Radical Beam

Detection is 1 m downstream of a nozzle.

Requirements for the Zeeman deceleration:

High-density, low rotational temperature, well collimated, slow,
and background free radical beam

UV dissociation of CH₃I strong background REMPI signal

UV dissociation or discharge of Di-tert-butyl peroxide

CH₃ Radical Beam

Detection is 1 m downstream of a nozzle.

Requirements for the Zeeman deceleration:

High-density, low rotational temperature, well collimated, slow, and background free radical beam

~~UV dissociation of CH₃I~~

strong background REMPI signal

UV dissociation or discharge of Di-tert-butyl peroxide

CH₃ Radical Beam

Detection is 1 m downstream of a nozzle.

Requirements for the Zeeman deceleration:

High-density, low rotational temperature, well collimated, slow,
and background free radical beam

~~UV dissociation of CH₃I~~

strong background REMPI signal

UV dissociation or discharge of Di-tert-butyl peroxide

high rotational temperature

CH₃ Radical Beam

Detection is 1 m downstream of a nozzle.

Requirements for the Zeeman deceleration:

High-density, low rotational temperature, well collimated, slow,
and background free radical beam

~~UV dissociation of CH₃I~~

strong background REMPI signal

~~UV dissociation or discharge of Di-tert-butyl peroxide~~

high rotational temperature

CH₃ Radical Beam

Detection is 1 m downstream of a nozzle.

Requirements for the Zeeman deceleration:

High-density, low rotational temperature, well collimated, slow, and background free radical beam


~~UV dissociation of CH₃I~~

strong background REMPI signal

~~UV dissociation or discharge of Di-tert-butyl peroxide~~

peroxide vs CH₄ discharge

high rotational temperature




Production of CH_3

Discharge of CH_4/Ar

Production of CH_3

Discharge of CH_4/Ar


(1) He seed gas is the best, but too fast.



Production of CH_3


Discharge of CH_4/Ar

(1) He seed gas is the best, but too fast.



(2) High-discharge voltage causes acceleration of radical beams


700 V



Production of CH₃

Discharge of CH₄/Ar


(1) He seed gas is the best, but too fast.





(2) High-discharge voltage causes acceleration of radical beams

700 V

480 V




(3) Cooling nozzle to 150 K makes the beam slow and intense




CH₃ signals are most intense at 150 K

Deceleration of CH_3

CH_4/Ar discharge




$1.25 \text{ cm}^{-1} / \text{coil}$




Phys. Chem. Chem. Phys **15**, 1772 (2013)

Deceleration of CH₃


CH₄/Ar discharge




1.25 cm⁻¹ /coil



15-coil system



expt.




sim.


Phys. Chem. Chem. Phys **15**, 1772 (2013)

Deceleration of CH₃

CH₄/Ar discharge



1.25 cm⁻¹ /coil





510 m/s → 470 m/s


$\Delta E \sim -19 \text{ cm}^{-1} \sim 28 \text{ K}$

Phys. Chem. Chem. Phys **15**, 1772 (2013)

15-coil system




Deceleration of CH₃ : 80 Coil System



Sample: 6.2bar 20%CH₄/Ar
Nozzle: 140K

Deceleration of CH₃ : 80 Coil System




Sample: 6.2bar 20%CH₄/Ar
Nozzle: 140K

Strong transverse focusing

CH₃ beam is less collimated



Deceleration of CH₃ : 80 Coil System



Sample: 6.2bar 20%CH₄/Ar
Nozzle: 140K


Strong transverse focusing

CH₃ beam is less collimated






Both $N = 0$ and 1 states are decelerated

Deceleration of CH₃ : 80 Coil System






Sample: 5bar 30%CH₄/Kr
Nozzle: 150K





Trapping of CH_3







Trapping of CH_3



Trapping of CH_3







Trapping of CH_3



CH_3 radicals are trapped !

Trapping of CH_3



CH_3 radicals are trapped !

Conclusions

CH₃ radicals are now spatially confined !

Anti-Helmholtz trap

$$v_{\parallel} = 40 \text{ ms}^{-1}$$

$$v_{\perp} \sim 1 \text{ ms}^{-1}$$



volume 4 mm³

Conclusions

CH₃ radicals are now spatially confined !

Anti-Helmholtz trap

$$v_{\parallel} = 40 \text{ ms}^{-1} \quad v_{\perp} \sim 1 \text{ ms}^{-1}$$



(1) high-resolution spectroscopy of CH₃

Doppler width < 500 kHz



volume 4 mm³

Conclusions

CH₃ radicals are now spatially confined !

Anti-Helmholtz trap

$$v_{\parallel} = 40 \text{ ms}^{-1} \quad v_{\perp} \sim 1 \text{ ms}^{-1}$$



volume 4 mm³

(1) high-resolution spectroscopy of CH₃

Doppler width < 500 kHz

(2) Cold collisions of CH₃

1 K ~ dB length 0.5 nm

Conclusions

CH₃ radicals are now spatially confined !

Anti-Helmholtz trap

$$v_{\parallel} = 40 \text{ ms}^{-1} \quad v_{\perp} \sim 1 \text{ ms}^{-1}$$



volume 4 mm³

(1) high-resolution spectroscopy of CH₃

Doppler width < 500 kHz

(2) Cold collisions of CH₃

1 K ~ dB length 0.5 nm

(3) Cold reactive collisions between CH₃ and O₂/OH

Conclusions

CH₃ radicals are now spatially confined !

Anti-Helmholtz trap

$$v_{\parallel} = 40 \text{ ms}^{-1} \quad v_{\perp} \sim 1 \text{ ms}^{-1}$$



volume 4 mm³

(1) high-resolution spectroscopy of CH₃

Doppler width < 500 kHz

(2) Cold collisions of CH₃

1 K ~ dB length 0.5 nm

(3) Cold reactive collisions between CH₃ and O₂/OH

(4) ortho-para conversion of CH₃

Conclusions

CH₃ radicals are now spatially confined !

Anti-Helmholtz trap



volume 4 mm³

$$v_{\parallel} = 40 \text{ ms}^{-1} \quad v_{\perp} \sim 1 \text{ ms}^{-1}$$

(1) high-resolution spectroscopy of CH₃

Doppler width < 500 kHz


(2) Cold collisions of CH₃

1 K ~ dB length 0.5 nm

(3) Cold reactive collisions between CH₃ and O₂/OH

(4) ortho-para conversion of CH₃

(5) sympathetic cooling with ultracold Rb/Li (<100 uK)



Conclusions

CH₃ radicals are now spatially confined !

Anti-Helmholtz trap



volume 4 mm³

$$v_{\parallel} = 40 \text{ ms}^{-1} \quad v_{\perp} \sim 1 \text{ ms}^{-1}$$

(1) high-resolution spectroscopy of CH₃

Doppler width < 500 kHz

(2) Cold collisions of CH₃


1 K ~ dB length 0.5 nm

(3) Cold reactive collisions between CH₃ and O₂/OH

(4) ortho-para conversion of CH₃

(5) sympathetic cooling with ultracold Rb/Li (<100 uK)

**Any non-singlet state of molecules
can be decelerated and trapped.**



Acknowledgements

Sida Zhou

Yang Liu

Pavle Djuricanin



Prof. Mark Raizen
(Texas Austin)



Dr. David Carty
(Durham, UK)

Cindy Toh (RI10)

Watheq Al-Basheer (RI12)



Tony Mittertreiner
(UBC/EES)



**NSERC
CRSNG**



Canada Foundation
for Innovation

Fondation canadienne
pour l'innovation

CRUCS

Canadian Centre
for Research on
Ultra-Cold Systems